



King's Research Portal

DOI:

[10.1016/j.archger.2018.03.011](https://doi.org/10.1016/j.archger.2018.03.011)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Li, D-J., Tseng, P-T., Stubbs, B., Chen, T-Y., Lin, P-Y., Chen, S-L., Thompson, T., Adamis, D., & Chu, C-S. (2018). Low peripheral levels of insulin growth factor-1 are associated with high incidence of delirium among elderly patients: A systematic review and meta-analysis. *ARCHIVES OF GERONTOLOGY AND GERIATRICS*. <https://doi.org/10.1016/j.archger.2018.03.011>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal


Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Title: Low peripheral levels of insulin growth factor-1 are associated with high incidence of delirium among elderly patients: A systematic review and meta-analysis

Authors: Dian-Jeng Li, Ping-Tao Tseng, Brendon Stubbs, Tien-Yu Chen, Pao-Yen Lin, Shiou-Lan Chen, Trevor Thompson, Dimitrios Adamis, Che-Sheng Chu



PII: S0167-4943(18)30047-5
DOI: <https://doi.org/10.1016/j.archger.2018.03.011>
Reference: AGG 3648

To appear in: *Archives of Gerontology and Geriatrics*

Received date: 21-12-2017
Revised date: 21-2-2018
Accepted date: 18-3-2018

Please cite this article as: Li, Dian-Jeng, Tseng, Ping-Tao, Stubbs, Brendon, Chen, Tien-Yu, Lin, Pao-Yen, Chen, Shiou-Lan, Thompson, Trevor, Adamis, Dimitrios, Chu, Che-Sheng, Low peripheral levels of insulin growth factor-1 are associated with high incidence of delirium among elderly patients: A systematic review and meta-analysis. *Archives of Gerontology and Geriatrics* <https://doi.org/10.1016/j.archger.2018.03.011>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Low peripheral levels of insulin growth factor-1 are associated with high incidence of delirium among elderly patients: A systematic review and meta-analysis

Running title: MA of IGF-1 and delirium

Dian-Jeng Li^{a,b,#}, Ping-Tao Tseng^{c,#}, Brendon Stubbs^{d,e,f}, Tien-Yu Chen^{g,h}, Pao-Yen Lin^{i,j},
Shiou-Lan Chen^b, Trevor Thompson^k, Dimitrios Adamis^{l,m}, Che-Sheng Chu^{n,o,*}

^a *Department of Addiction Science, Kaohsiung Municipal Kai-Syuan Psychiatric Hospital, Kaohsiung City, Taiwan*

^b *Graduate Institute of Medicine, College of Medicine, Kaohsiung Medical University, Taiwan*

^c *WinShine Clinics in Specialty of Psychiatry, Kaohsiung City, Taiwan*

^d *Physiotherapy Department, South London and Maudsley NHS Foundation Trust, London, UK*

^e *Health Service and Population Research Department, Institute of Psychiatry, Psychology and Neuroscience (IoPPN), King's College London, De Crespigny Park, London, UK*

^f *Faculty of Health, Social Care and Education, Anglia Ruskin University, Chelmsford, UK*

^g *Department of Psychiatry, Tri-Service General Hospital; School of Medicine, National Defense Medical Center, Taipei, Taiwan*

^h *Institute of Brain Science, National Yang-Ming University, Taipei, Taiwan*

ⁱ *Department of Psychiatry, Kaohsiung Chang Gung Memorial Hospital and Chang Gung University College of Medicine, Kaohsiung, Taiwan*

^j *Institute for Translational Research in Biomedical Sciences, Kaohsiung Chang Gung Memorial Hospital*

^k *Faculty of Education and Health, University of Greenwich, London SE9 2UG*

^l *Sligo Mental Health Services Clarion Rd Sligo, Ireland*

^m *Research and Academic Institute of Athens, Greece*

ⁿ *Department of Psychiatry, Kaohsiung Veterans General Hospital, Kaohsiung, Taiwan*

^o *Center for Geriatric and Gerontology, Kaohsiung Veterans General Hospital, Kaohsiung, Taiwan*

[#] *contributed equally as first author*

*** Correspondence to:**

Che-Sheng Chu, MD

Department of Psychiatry, Kaohsiung Veterans General Hospital, Kaohsiung, Taiwan

Address: No. 386, Dazhong 1st Rd., Zuoying Dist., Kaohsiung City 81362, Taiwan

Telephone: 886-7-3422121

Fax: 886-7-3422288

E-mail: youngtzuchi@hotmail.com/cschu@vghks.gov.tw

Highlights

- The study provides a comprehensive and quantitative review of IGF-1 and delirium.
- IGF-1 might play a role in the pathogenesis in the delirium.
- Lower peripheral levels of IGF-1 could be associated with delirium in the elderly.

Abstract:

Introduction: Delirium, a serious condition observed in critically ill patients, clinically presents with impaired cognition and consciousness. The relationship between delirium and peripheral levels of insulin growth factor-1 (IGF-1) is unclear. Thus we conducted a meta-analysis to address this issue.

Methods: Seven major electronic databases were searched from inception until October 2, 2017

to obtain relevant clinical variables to compare the difference in IGF-1 levels between delirious and non-delirious elderly in-patients. A random effects meta-analysis was conducted.

Results: We studies 10 articles involving 294 delirious patients (mean age 73.0 years) and 604 non-delirious patients (mean age 76.9 years). We found that peripheral levels of IGF-1 in patients with delirium were significantly lower than in those without delirium (Hedges' $g = -0.209$, 95% confidence interval [CI] = -0.393 to -0.026, $p = 0.025$). Meta-regression analyses found that no variables such as percentage of cognitive impairment, mean age, and female proportion contribute to heterogeneity in terms of the entire population.

Conclusions: Our data suggests that lower peripheral levels of IGF-1 could be associated with a higher incidence of delirium among elderly patients. Further prospective studies with larger sample sizes are needed to investigate the association between peripheral levels of IGF-1 and delirium.

Abbreviation: CI: confidence interval; IGF-1: insulin-like growth factor-1

Keywords: Biomarker; Delirium; Elderly; IGF-1; Meta-analysis

1. Introduction

Delirium, which clinically presents with impairment in cognition, consciousness, psychomotor behavior, and attention (Ely et al., 2001), may manifest with a wide range of subtypes ranging between hyperactive and hypoactive conditions (Steiner, 2011). Clinically, it usually presents with a sudden onset, which then fluctuates between hours and days and is generally a reversible condition (Yudofsky & Hales, 2008). Etiologically, delirium indicates an organic decline in cerebral function resulting from variable mechanisms. For example, the etiology of alcohol-withdrawal delirium differs from that causing postoperative delirium (Steiner, 2011). Incidence of delirium varies depending on the purpose of investigation. Incidence of delirium varies between 9% and 43% among geriatric patients hospitalized for variable medical conditions (Laurila, Pitkala, Strandberg, & Tilvis, 2002; O'Keeffe & Gosney, 1997). Delirium is commonly observed in elderly patients post surgery, and the incidence of postoperative delirium (POD) ranges widely between 10% and 65% (Radtke et al., 2010; Rudolph & Marcantonio, 2011). Delirium is a serious neuropsychiatric condition with a substantial impact owing to increased morbidity and mortality, high healthcare costs and prolonged hospitalization associated with it (Hustey, Meldon, Smith, & Lex, 2003;

Meagher et al., 2007). Regarding mortality rates, reportedly, delirious patients show an overall hazard ratio of 1.9 in elderly patients in a meta-analysis study (Witlox et al., 2010). Given the inadequate understanding of its multifactorial aetiopathogenesis (Hughes, Patel, & Pandharipande, 2012), researchers are interested in studying the etiology of delirium to better predict its course and prognosis in clinical settings. Although previous reviews (Khan, Zawahiri, Campbell, & Boustani, 2011) have demonstrated the limited role of biomarkers in detection of delirium, researchers have been trying to identify specific biomarkers linked to this condition such as insulin growth factor-1 (IGF-1).

IGF-1 is a member of the IGF family with 70 amino acid peptides, and is primarily produced in the central nervous system (CNS) and peripheral tissues such as the liver (Le Roith, 2003; Torres-Aleman, 2010). IGF-1 in the CNS plays a vital role in enhancing neuronal survival, differentiation, synaptogenesis, and recovery from neural injury (D'Ercole, Ye, & O'Kusky, 2002). IGF-I blocks apoptosis in damaged neurons via several mechanisms, such as delay of apoptosis via activation of PI3K/Akt kinase and Ras/Raf/MEK/ERK pathways and results in long-term survival (Gluckman et al., 1998; Parrizas & LeRoith, 1997; Peruzzi et al., 1999). Clinically, studies have explored its neuroprotective role and association with neurocognitive disorders such as Alzheimer's disease (AD), of which is involved with delirium (Kooijman, Sarre, Michotte, & De Keyser, 2009; Lackey, Gray, & Henricks, 2000). Additionally, IGF-1 deficiency in humans may cause growth abnormalities, including a

smaller sized brain and mental retardation (Werner & LeRoith, 2014). Furthermore, it has been reported that serum IGF-1 levels positively correlate with cognitive performance in healthy elderly subjects (Al-Delaimy, von Muhlen, & Barrett-Connor, 2009; Aleman et al., 2000).

Moreover, as researchers keenly investigating the association between peripheral levels of IGF-1 and delirium, several studies have explored the significance of this correlation in patients undergoing surgery or those with acute medical illnesses (Adamis et al., 2009; Adamis et al., 2007; Cerejeira, Batista, Nogueira, Vaz-Serra, & Mukaetova-Ladinska, 2013; Egberts et al., 2015; Lemstra, Kalisvaart, Vreeswijk, van Gool, & Eikelenboom, 2008; Morandi et al., 2011; Wilson, Broadhurst, Diver, Jackson, & Mottram, 2005). While some studies report that lower peripheral levels of IGF-I could significantly increase the incidence of delirium (Cerejeira et al., 2013; Egberts et al., 2015; Morandi et al., 2011; Wilson et al., 2005), the opposite findings have also been reported by some others (Adamis et al., 2009; Adamis et al., 2007; Lemstra et al., 2008). Possible factors accounting for this discrepancy are inconsistency between recruited patients, such as postoperative or medically ill cases in terms of study designs or clinical status. Thus, the association between peripheral levels of IGF-I and incidence of delirium in a clinical setting needs to be further studied. Given these aforementioned gaps existing in literature, we performed a comprehensive systematic review and meta-analysis investigating the association between peripheral levels of IGF-1 and

delirium, specifically focusing on clinical status of patients. In addition to exploring the difference in this association between surgical or medically ill patients, we also attempted to determine possible factors affecting the correlation between delirium and IGF-1 levels.

2. Materials and methods

Our systematic review and meta-analysis was based on Meta-analysis Of Observational Studies in Epidemiology (MOOSE) guidelines (Stroup et al., 2000) (supplemental table S1) and was approved by the Institutional Review Board of the Tri-Service General Hospital (TSGHIRB: B-105-12). Two investigators (DJ Li and CS Chu) independently performed database searches, study selection, data extraction, and rating of the methodological quality of included studies. Disagreements between investigators were resolved through consensus.

2.1. Database searches

PubMed, Embase, ScienceDirect, ClinicalKey, Cochrane Library, ClinicalTrials.gov and ProQuest were systematically searched from the time of their inception until October 2nd, 2017. Keywords used for our search included “(IGF OR IGF1 protein OR inflammation mediators OR insulin-like growth factor OR cytokine) AND (delirium OR acute confusion OR consciousness disturbance)” without any limitation about language. Detailed search strings used for the electronic database search and results are shown in the supplemental figure S1.

2.2. Eligibility Criteria and Study Selection

Eligibility criteria applied were: (1) case-control studies comparing peripheral IGF-1 levels between delirious patients and non-delirious controls, or cohorts with detailed baseline data and subsequent delirium risk; (2) those with a diagnosis of delirium based on the definite diagnostic systems or scales, such as confusion assessment method (CAM) (Inouye et al., 1990), the diagnostic and statistical manual of mental disorders, 3rd revision (American Psychiatric Association. 1980) and text-revision (DSM-IV-TR) (American Psychiatric Association. 2000), or Delirium observation screening scale (Schuurmans, Shortridge-Baggett, & Duursma, 2003), Richmond agitation-sedation scale (Sessler et al., 2002); (3) those in whom detailed methods for detecting peripheral IGF-1 were available; and (4) formal registered trials or formal published peer-reviewed articles.

Exclusion criteria were: (1) studies without adequate non-delirious controls; (2) articles without relation to the comparisons of peripheral IGF-1 levels in delirious patients and controls; (3) those irrelevant to delirium; (4) review articles/commentary articles/case reports/letter to editor; or (5) non-clinical trials in human.

2.3. Methodological Quality Assessment

The modified Newcastle-Ottawa Scale (NOS) was used to rate methodological quality of included studies (Wells GA). This scale considers three domains: selection of participants

(maximum of four stars), comparability of groups (maximum of two stars), and measurement (maximum of three stars). Scores ranged from 0 (the lowest) to 9 (the highest), and a score equal to 7 indicated high methodological quality.

2.4. Data Extraction

A predetermined data extraction list was used to extract data for this meta-analysis. To be specific, the extracted data included basic characteristics of participants (mean age and percentage of female), mini-mental status examination (MMSE), and duration of hospitalization.

2.5. Meta-analysis

The primary outcomes were the difference in peripheral IGF-1 levels in delirious patients compared to control groups who did not develop delirium in each recruited study (calculated as Hedges' g statistic). We did not choose difference in means as effect sizes (ESs) of our primary outcome due to presumed different units used among each study. For studies in which peripheral levels of IGF-1 were not available, outcomes were chosen as odd ratios (ORs) (Morandi et al., 2011; Wilson et al., 2005), or only a p value (Adamis et al., 2007) according to manuals of Comprehensive Meta-Analysis version 3.0. When peripheral levels of IGF-1 were not available in the included studies, we tried to contact the authors of such studies on at least two separate occasions to try and obtain additional raw data. Additionally,

we converted ORs into ES using the formula suggested by the Cochrane Handbook (Higgins & Green, 2011) and derived ES using other statistical parameters such as the t or p value with the sample size when raw data were not available via email inquiry. We defined an ES and Hedges' g less than 0 as "lower peripheral IGF-1 levels in patients with delirium than in controls without delirium". Statistical significance was set as a two-tailed p value < 0.05 . All meta-analytical procedures were performed using the Comprehensive Meta-Analysis version 3 (CMA ver. 3.0) (Biostat, Englewood, NJ) software. Meta-analyses were performed using random-effects models due to the anticipated heterogeneity across studies. Briefly, random-effects modeling is more stringent and therefore more convincing than fixed-effects modeling and incorporates a between-study variance in the calculations (Borenstein, Hedges, Higgins, & Rothstein, 2010).

2.6. *Heterogeneity, publication bias, sensitivity test, and meta-regression*

We used the Cochran's Q test and the corresponding p value to investigate potential heterogeneity among recruited studies (Higgins & Thompson, 2002). The I^2 metric was used to investigate the proportion of variance between recruited studies (Borenstein, Higgins, Hedges, & Rothstein, 2017). The publication bias was determined by visual examination of funnel plots in situations of less than 10 studies (Higgins & Green, 2011) and using thorough Egger's regression analysis if there were more than 10 studies (Egger, Davey Smith,

Schneider, & Minder, 1997). In situation of significant publication bias, we arranged Duval and Tweedie's trim and fill test to adjust the results of meta-analysis (Duval & Tweedie, 2000). Additionally, we performed sensitivity testing with one study removal test to investigate potential confounding by any one of the outlier in the recruited studies, which had been widely used in meta-analysis (Tobias, 1999). Additionally, we performed meta-regression analysis using an unrestricted maximum likelihood method to explore potential moderators. Variables considered for meta-regression analyses were: mean age, female gender (%), MMSE score, and a total modified NOS scores. The meta-regression procedure was only performed when moderator variables were available from more than 5 individual datasets.

3. Results

3.1. Studies included in the meta-analysis

The whole search picture was depicted as figure 1 and supplemental figure S1. In brief, eighty-two articles enter the full-texts investigation stage, and seventy-two were excluded for specific reasons (see supplemental table S2). Finally, ten articles met our inclusion criteria (Adamis et al., 2007; Cerejeira et al., 2013; Chu et al., 2016; Cinar et al., 2014; Egberts et al., 2015; Lemstra et al., 2008; Morandi et al., 2011; Shen, Shao, Chen, & Guo, 2016; Wilson et al., 2005; Yen et al., 2016). Peripheral levels of IGF-1 were obtained before delirium

occurring in the seven studies (Cerejeira et al., 2013; Chu et al., 2016; Cinar et al., 2014; Lemstra et al., 2008; Shen et al., 2016; Wilson et al., 2005; Yen et al., 2016). The participants in other three studies might be drawn IGF-1 during delirious status (Adamis et al., 2007; Egberts et al., 2015; Morandi et al., 2011).

3.2. Methodological quality assessment

The mean modified NOS score was 7.4 with standard deviation (SD) = 0.5 across the ten studies. The score in each domain of the modified NOS is shown in supplemental table S3.

3.3. The main results of the meta-analysis comparing the levels of IGF-1 in patients with delirium and those without delirium

Our systematic review and meta-analysis included 10 articles that met our inclusion criteria. Peripheral (serum or plasma) IGF-1 levels were measured in these ten studies. The quantitative determination of IGF-1 were taken from plasma (Cerejeira et al., 2013; Wilson et al., 2005) and serum (Adamis et al., 2007; Chu et al., 2016; Cinar et al., 2014; Egberts et al., 2015; Morandi et al., 2011; Shen et al., 2016; Yen et al., 2016), respectively. One study (Lemstra et al., 2008) mentioned that IGF-1 was assessed from blood sample (table 1). We found 294 delirious patients (mean age 73.0 years, women 51.0 %) and 604 non-delirious patients (mean age 76.9 years, women 53.2 %) among the 10 studies. Peripheral levels of IGF-1 in patients with delirium were significantly lower than in those without delirium

(Hedges' $g = -0.209$, 95% confidence interval (CI) = -0.393 to -0.026, $p = 0.025$) (Figure 2)

with significant heterogeneity (Q value = 20.215, degree of freedom (df) = 9, $I^2 = 55.479\%$, $p = 0.017$) but no significant publication bias based on Egger's test [t value = 1.060, df = 8, $p = 0.320$].

3.3.1. Sensitivity test

The significant result of current meta-analysis became non-significant after removal of the datasets by Adamis, D. (2007) (Hedges' $g = -0.184$, 95% CI = -0.374 to 0.006, $p = 0.058$) (Adamis et al., 2007), Egberts, A. (2015) (Hedges' $g = -0.175$, 95% CI = -0.362 to 0.011, $p = 0.066$) (Egberts et al., 2015), Shen, H. (2016) (Hedges' $g = -0.125$, 95% CI = -0.262 to 0.012, $p = 0.073$) (Shen et al., 2016), or Yen, T.E. (2016) (Hedges' $g = -0.193$, 95% CI = -0.390 to 0.004, $p = 0.055$) (Yen et al., 2016).

3.3.2 Meta-regression

There was no any significant association between the differences of peripheral IGF-1 and clinical variables, including percentage of cognitive impairment ($p=0.960$), mean age ($p=0.978$), female proportion ($p=0.721$), mean MMSE scores ($p=0.929$), or quality of methodology as total modified NOS scores ($p=0.270$).

4. Discussion

To the best of our knowledge, the study is the first meta-analysis to address the association between peripheral IGF-1 and delirium and suggests that lower peripheral IGF-1 levels were found in patients with delirium compared with those without delirium. Although causality of delirium and peripheral levels of IGF-1 has yet to be clarified, our study offers additional evidence supporting the immune-inflammatory hypothesis in delirium.

The link between delirium and peripheral levels of IGF-1 can be mediated by cytokines released during inflammatory processes. Reportedly, a strong correlation is seen to exist between delirium and inflammation (Broadhurst & Wilson, 2001). Peripheral IGF-1 can cross the blood-brain barrier and play a role in neuronal function (Reinhardt & Bondy, 1994). It has been reported that administration of IGF-1 reduces secondary neuronal loss after transient neural injuries (Gluckman et al., 1998; Trejo, Carro, Garcia-Galloway, & Torres-Aleman, 2004). Similarly, replacement therapy with IGF-1 ameliorates diabetic neuropathy (Schmidt et al., 1999). Our study provides additional evidence to support this hypothesis. Lower peripheral levels of IGF-1 may possess lesser neuroprotective properties that is anti-apoptotic and anti-inflammatory actions, and contributed higher risk of delirium.

Another factor involved in the connection between peripheral levels of IGF-1 and delirium may be relatively high mean age (almost above 70 years old) of participants in the included studies. Ageing, resulting from increased natural immunity activity, is a low level

inflammatory status (Godbout & Johnson, 2009; Licastro et al., 2005). Inflammation is related to several morbid conditions in the elderly such as cardiovascular disease (Pai et al., 2004) or Alzheimer's disease (Cassidy & Topol, 2004). Furthermore, inflammation may result in the breakdown of the blood-brain barrier (Jeppsson et al., 1981) and decrease cholinergic transfer (Hammond, Meador, Aung-Din, & Wilder, 1987). Taken together, elevated levels of proinflammatory cytokine levels have been suggested to predispose the development of delirium in elderly, which is interpreted by the concept of immunosenescence (Godbout & Johnson, 2006). IGF-1, involving neuro-inflammation in the CNS (Trejo et al., 2004), may be also affected by ageing, be associated with delirium. However, our meta-regression showed that mean age did not significantly involve the relation between delirium and IGF-1. In addition to limited case numbers, relatively narrowing distribution of mean age may also restrict the statistical significance. Previous study also indicated that older patients undergoing cardiovascular surgery may develop delirium easily (Cinar et al., 2014). However, clearly more research is required to better understand this issue.

It has been reported that IGF-1 levels positively correlate with cognitive performance in healthy elderly subjects (Al-Delaimy et al., 2009; Aleman et al., 2000). A meta-analysis has reported a significantly positive correlation between IGF-I and cognition in a healthy elderly population (Arwert, Deijen, & Drent, 2005). Another study concluded that the IGFI levels showed an inverse correlation with cognitive dysfunction (Murialdo et al., 2001). In the

present meta-analysis study, only two studies enrolled participants with baseline cognitive healthy, which was defined as an MMSE score of ≥ 24 or excluded when enrollment, and reported that lower peripheral levels of IGF-1 associated with incident of delirium in one study (adjusted Odd ratios [aORs] = 2.52, 95% CI = 1.19 to 5.43, $p = 0.019$) (Shen et al., 2016) but not another (aORs = 0.99, 95% CI = 0.96 to 1.01, $p = 0.183$) (Yen et al., 2016). For participants with baseline cognitive impairment, six (Adamis et al., 2007; Cerejeira et al., 2013; Chu et al., 2016; Cinar et al., 2014; Lemstra et al., 2008; Morandi et al., 2011) out of eight (Adamis et al., 2007; Cerejeira et al., 2013; Chu et al., 2016; Cinar et al., 2014; Egberts et al., 2015; Lemstra et al., 2008; Morandi et al., 2011; Wilson et al., 2005) studies showed no significant differences between peripheral levels of IGF-1 and delirium. Therefore, cognitive impairment could be a possible confounder unfavorably affecting the association between delirium and peripheral levels of IGF-1. However, our meta-regression showed that baseline MMSE scores were not significantly linked to a correlation between delirium and IGF-1. This difference may be due to variable statistical methodology between dichotomous grouping [normal cognition vs. mild cognitive impairment (MCI) or dementia] and continuous variables of baseline MMSE scores. The fact that a limited number of cases were studied might contribute to the insignificance observed with respect to baseline MMSE scores.

Furthermore, some studies report that IGF-I levels were unaltered or increased among dementia patients (Nasman et al., 1995; Sara et al., 1982; Tham et al., 1993). A greater

number of studies will need to be performed in future to investigate the correlation between cognition and IGF-1 to explore this inconsistency.

Limitations of our study that need to be addressed are: 1) Studying a limited number of cases might be a confounding factor impacting the significance of outcomes. 2) Higher mean age would restrict the application of our findings to younger adults or adolescents. 3) We could not perform subgroup analyses comparing IGF-1 levels in different delirious classifications or diagnostic criteria (hypoactive vs hyperactive delirium and only CAM vs CAM plus DSM-III or -IV diagnosis) due to not enough studies. Future research is required to understand if this influences the relationship observed. 4) Peripheral levels of IGF-1 do not actually reflect the brain levels, and thus the pathogenesis between CNS levels of IGF-1 and incident delirium need to be addressed in future studies.

5. Conclusion

In conclusion, lower peripheral levels of IGF-1 appear to be associated with delirium among elderly patients. Further studies involving a greater number of participants and wider ranges of age would help to investigate the etiology and association between these two entities in future. Studies including more baseline variables and more specific description of diagnosing method to delirium could help us investigate more covariates and conduct subgroup analysis.

Conflict of Interest

The authors state that there are no any competing interests in the current literature.

Acknowledgements

The authors declare that there are no conflicts of interest or funding in relation to the subject of this study.

References:

- Adamis, D., Lunn, M., Martin, F. C., Treloar, A., Gregson, N., Hamilton, G., & Macdonald, A. J. (2009). Cytokines and IGF-I in delirious and non-delirious acutely ill older medical inpatients. *Age Ageing*, 38(3), 326-332.
- Adamis, D., Treloar, A., Martin, F. C., Gregson, N., Hamilton, G., & Macdonald, A. J. (2007). APOE and cytokines as biological markers for recovery of prevalent delirium in elderly medical inpatients. *Int J Geriatr Psychiatry*, 22(7), 688-694.
- Al-Delaimy, W. K., von Muhlen, D., & Barrett-Connor, E. (2009). Insulinlike growth factor-1, insulinlike growth factor binding protein-1, and cognitive function in older men and women. *J Am Geriatr Soc*, 57(8), 1441-1446.
- Aleman, A., de Vries, W. R., de Haan, E. H., Verhaar, H. J., Samson, M. M., & Koppeschaar, H. P. (2000). Age-sensitive cognitive function, growth hormone and insulin-like growth factor 1 plasma levels in healthy older men. *Neuropsychobiology*, 41(2), 73-78.
- American Psychiatric Association. (1980). *Diagnostic and Statistical Manual of Mental Disorder* (3rd ed., text rev.). Washington, DC: American Psychiatric Association.
- American Psychiatric Association. (2000). *Diagnostic and Statistical Manual of*

Mental Disorder (4th ed., text rev.). Washington, DC: American Psychiatric Association.

Arwert, L. I., Deijen, J. B., & Drent, M. L. (2005). The relation between insulin-like growth factor I levels and cognition in healthy elderly: a meta-analysis.

Growth Horm IGF Res, 15(6), 416-422.

Borenstein, M., Hedges, L. V., Higgins, J. P., & Rothstein, H. R. (2010). A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res Synth Methods*, 1(2), 97-111.

Borenstein, M., Higgins, J. P., Hedges, L. V., & Rothstein, H. R. (2017). Basics of meta-analysis: I² is not an absolute measure of heterogeneity. *Res Synth Methods*, 8(1), 5-18.

Broadhurst, C., & Wilson, K. (2001). Immunology of delirium: new opportunities for treatment and research. *Br J Psychiatry*, 179, 288-289.

Casserly, I., & Topol, E. (2004). Convergence of atherosclerosis and Alzheimer's disease: inflammation, cholesterol, and misfolded proteins. *Lancet*, 363(9415), 1139-1146.

Cerejeira, J., Batista, P., Nogueira, V., Vaz-Serra, A., & Mukaetova-Ladinska, E. B.

(2013). The stress response to surgery and postoperative delirium: evidence of hypothalamic-pituitary-adrenal axis hyperresponsiveness and decreased

suppression of the GH/IGF-1 Axis. *J Geriatr Psychiatry Neurol*, 26(3), 185-194.

Chu, C. S., Liang, C. K., Chou, M. Y., Lin, Y. T., Hsu, C. J., Chu, C. L., & Chou, P.

H. (2016). Lack of Association between Pre-Operative Insulin-Like Growth Factor-1 and the Risk of Post-Operative Delirium in Elderly Chinese Patients. *Psychiatry Investig*, 13(3), 327-332.

Cinar, M. A., Balikci, A., Sertoglu, E., Mehmet, A. K., Serdar, M. A., & Ozmenler,

K. N. (2014). Role of CRP, TNF- α , and IGF-1 in Delirium Pathophysiology. *Noro Psikiyatr Ars*, 51(4), 376-382.

D'Ercole, A. J., Ye, P., & O'Kusky, J. R. (2002). Mutant mouse models of insulin-like growth factor actions in the central nervous system. *Neuropeptides*, 36(2-3), 209-220.

Duval, S., & Tweedie, R. (2000). Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, 56(2), 455-463.

Egberts, A., Wijnbeld, E. H., Fekkes, D., van der Ploeg, M. A., Ziere, G., Hooijkaas,

H., . . . Mattace-Raso, F. U. (2015). Neopterin: a potential biomarker for delirium in elderly patients. *Dement Geriatr Cogn Disord*, 39(1-2), 116-124.

Egger, M., Davey Smith, G., Schneider, M., & Minder, C. (1997). Bias in meta-

analysis detected by a simple, graphical test. *BMJ*, 315(7109), 629-634.

Ely, E. W., Inouye, S. K., Bernard, G. R., Gordon, S., Francis, J., May, L., . . . Dittus,

R. (2001). Delirium in mechanically ventilated patients: validity and reliability of the confusion assessment method for the intensive care unit (CAM-ICU).

JAMA, 286(21), 2703-2710.

Gluckman, P. D., Guan, J., Williams, C., Scheepens, A., Zhang, R., Bennet, L., &

Gunn, A. (1998). Asphyxial brain injury--the role of the IGF system. *Mol Cell Endocrinol*, 140(1-2), 95-99.

Godbout, J. P., & Johnson, R. W. (2006). Age and neuroinflammation: a lifetime of psychoneuroimmune consequences. *Neurol Clin*, 24(3), 521-538.

Godbout, J. P., & Johnson, R. W. (2009). Age and neuroinflammation: a lifetime of psychoneuroimmune consequences. *Immunol Allergy Clin North Am*, 29(2), 321-337.

Hammond, E. J., Meador, K. J., Aung-Din, R., & Wilder, B. J. (1987). Cholinergic modulation of human P3 event-related potentials. *Neurology*, 37(2), 346-350.

Higgins, J. P., & Green, S. (2011). *Cochrane Handbook for Systematic Reviews of Interventions* (5.1.0 ed.): The Cochrane Collaboration. Available from www.handbook.cochrane.org.

Higgins, J. P., & Thompson, S. G. (2002). Quantifying heterogeneity in a meta-

analysis. *Stat Med*, 21(11), 1539-1558.

Hughes, C. G., Patel, M. B., & Pandharipande, P. P. (2012). Pathophysiology of acute brain dysfunction: what's the cause of all this confusion? *Curr Opin Crit Care*, 18(5), 518-526.

Hustey, F. M., Meldon, S. W., Smith, M. D., & Lex, C. K. (2003). The effect of mental status screening on the care of elderly emergency department patients. *Ann Emerg Med*, 41(5), 678-684.

Inouye, S. K., van Dyck, C. H., Alessi, C. A., Balkin, S., Siegel, A. P., & Horwitz, R. I. (1990). Clarifying confusion: the confusion assessment method. A new method for detection of delirium. *Ann Intern Med*, 113(12), 941-948.

Jeppsson, B., Freund, H. R., Gimmon, Z., James, J. H., von Meyenfeldt, M. F., & Fischer, J. E. (1981). Blood-brain barrier derangement in sepsis: cause of septic encephalopathy? *Am J Surg*, 141(1), 136-142.

Khan, B. A., Zawahiri, M., Campbell, N. L., & Boustani, M. A. (2011). Biomarkers for delirium--a review. *J Am Geriatr Soc*, 59 Suppl 2, S256-261.

Kooijman, R., Sarre, S., Michotte, Y., & De Keyser, J. (2009). Insulin-like growth factor I: a potential neuroprotective compound for the treatment of acute ischemic stroke? *Stroke*, 40(4), e83-88.

Lackey, B. R., Gray, S. L., & Henricks, D. M. (2000). Actions and interactions of the

IGF system in Alzheimer's disease: review and hypotheses. *Growth Horm IGF Res*, 10(1), 1-13.

Laurila, J. V., Pitkala, K. H., Strandberg, T. E., & Tilvis, R. S. (2002). Confusion assessment method in the diagnostics of delirium among aged hospital patients: would it serve better in screening than as a diagnostic instrument? *Int J Geriatr Psychiatry*, 17(12), 1112-1119.

Le Roith, D. (2003). The insulin-like growth factor system. *Exp Diabesity Res*, 4(4), 205-212.

Lemstra, A. W., Kalisvaart, K. J., Vreeswijk, R., van Gool, W. A., & Eikelenboom, P. (2008). Pre-operative inflammatory markers and the risk of postoperative delirium in elderly patients. *Int J Geriatr Psychiatry*, 23(9), 943-948.

Licastro, F., Candore, G., Lio, D., Porcellini, E., Colonna-Romano, G., Franceschi, C., & Caruso, C. (2005). Innate immunity and inflammation in ageing: a key for understanding age-related diseases. *Immun Ageing*, 2, 8.

Meagher, D. J., Moran, M., Raju, B., Gibbons, D., Donnelly, S., Saunders, J., & Trzepacz, P. T. (2007). Phenomenology of delirium. Assessment of 100 adult cases using standardised measures. *Br J Psychiatry*, 190, 135-141.

Morandi, A., Gunther, M. L., Pandharipande, P. P., Jackson, J. C., Thompson, J. L., Shintani, A. K., . . . Girard, T. D. (2011). Insulin-like growth factor-1 and

delirium in critically ill mechanically ventilated patients: a preliminary investigation. *Int Psychogeriatr*, 23(7), 1175-1181.

Murialdo, G., Barreca, A., Nobili, F., Rollero, A., Timossi, G., Gianelli, M. V., . . .

Polleri, A. (2001). Relationships between cortisol, dehydroepiandrosterone sulphate and insulin-like growth factor-I system in dementia. *J Endocrinol Invest*, 24(3), 139-146.

Nasman, B., Olsson, T., Seckl, J. R., Eriksson, S., Viitanen, M., Bucht, G., &

Carlstrom, K. (1995). Abnormalities in adrenal androgens, but not of glucocorticoids, in early Alzheimer's disease. *Psychoneuroendocrinology*, 20(1), 83-94.

O'Keeffe, S. T., & Gosney, M. A. (1997). Assessing attentiveness in older hospital patients: global assessment versus tests of attention. *J Am Geriatr Soc*, 45(4), 470-473.

Pai, J. K., Pischon, T., Ma, J., Manson, J. E., Hankinson, S. E., Joshipura, K., . . .

Rimm, E. B. (2004). Inflammatory markers and the risk of coronary heart disease in men and women. *N Engl J Med*, 351(25), 2599-2610.

Parrizas, M., & LeRoith, D. (1997). Insulin-like growth factor-1 inhibition of

apoptosis is associated with increased expression of the bcl-xL gene product. *Endocrinology*, 138(3), 1355-1358.

Peruzzi, F., Prisco, M., Dews, M., Salomoni, P., Grassilli, E., Romano, G., . . .

Baserga, R. (1999). Multiple signaling pathways of the insulin-like growth factor 1 receptor in protection from apoptosis. *Mol Cell Biol*, 19(10), 7203-7215.

Radtke, F. M., Franck, M., MacGuill, M., Seeling, M., Lutz, A., Westhoff, S., . . .

Spies, C. D. (2010). Duration of fluid fasting and choice of analgesic are modifiable factors for early postoperative delirium. *Eur J Anaesthesiol*, 27(5), 411-416.

Reinhardt, R. R., & Bondy, C. A. (1994). Insulin-like growth factors cross the blood-brain barrier. *Endocrinology*, 135(5), 1753-1761.

Rudolph, J. L., & Marcantonio, E. R. (2011). Review articles: postoperative delirium: acute change with long-term implications. *Anesth Analg*, 112(5), 1202-1211.

Sara, V. R., Hall, K., Enzell, K., Gardner, A., Morawski, R., & Wetterberg, L. (1982).

Somatomedins in aging and dementia disorders of the Alzheimer type. *Neurobiol Aging*, 3(2), 117-120.

Schmidt, R. E., Dorsey, D. A., Beaudet, L. N., Plurad, S. B., Parvin, C. A., & Miller,

M. S. (1999). Insulin-like growth factor I reverses experimental diabetic autonomic neuropathy. *Am J Pathol*, 155(5), 1651-1660.

Schuurmans, M. J., Shortridge-Baggett, L. M., & Duursma, S. A. (2003). The

Delirium Observation Screening Scale: a screening instrument for delirium.

Res Theory Nurs Pract, 17(1), 31-50.

Sessler, C. N., Gosnell, M. S., Grap, M. J., Brophy, G. M., O'Neal, P. V., Keane, K.

A., . . . Elswick, R. K. (2002). The Richmond Agitation-Sedation Scale:

validity and reliability in adult intensive care unit patients. *Am J Respir Crit*

Care Med, 166(10), 1338-1344.

Shen, H., Shao, Y., Chen, J., & Guo, J. (2016). Insulin-Like Growth Factor-1, a

Potential Predictive Biomarker for Postoperative Delirium Among Elderly

Patients with Open Abdominal Surgery. *Curr Pharm Des*, 22(38), 5879-5883.

Steiner, L. A. (2011). Postoperative delirium. Part 1: pathophysiology and risk

factors. *Eur J Anaesthesiol*, 28(9), 628-636.

Stroup, D. F., Berlin, J. A., Morton, S. C., Olkin, I., Williamson, G. D., Rennie, D., . .

. Thacker, S. B. (2000). Meta-analysis of observational studies in

epidemiology: a proposal for reporting. Meta-analysis Of Observational

Studies in Epidemiology (MOOSE) group. *JAMA*, 283(15), 2008-2012.

Tham, A., Nordberg, A., Grissom, F. E., Carlsson-Skwirut, C., Viitanen, M., & Sara,

V. R. (1993). Insulin-like growth factors and insulin-like growth factor

binding proteins in cerebrospinal fluid and serum of patients with dementia of

the Alzheimer type. *J Neural Transm Park Dis Dement Sect*, 5(3), 165-176.

Tobias, A. (1999). Assessing the influence of a single study in meta-analysis. *STATA*

technical bulletin, 47, 15–17.

Torres-Aleman, I. (2010). Toward a comprehensive neurobiology of IGF-I. *Dev*

Neurobiol, 70(5), 384-396.

Trejo, J. L., Carro, E., Garcia-Galloway, E., & Torres-Aleman, I. (2004). Role of

insulin-like growth factor I signaling in neurodegenerative diseases. *J Mol*

Med (Berl), 82(3), 156-162.

Wells GA, S. B., O'Connell D, Peterson J, Welch V, et al.. (2011) The Newcastle-

Ottawa Scale (NOS) for assessing the quality of nonrandomized studies in

meta-analysis. Available:

www.ohri.ca/programs/clinical_epidemiology/oxford.asp.

Werner, H., & LeRoith, D. (2014). Insulin and insulin-like growth factor receptors in

the brain: physiological and pathological aspects. *Eur Neuropsychopharmacol*,

24(12), 1947-1953.

Wilson, K., Broadhurst, C., Diver, M., Jackson, M., & Mottram, P. (2005). Plasma

insulin growth factor-1 and incident delirium in older people. *Int J Geriatr*

Psychiatry, 20(2), 154-159.

Witlox, J., Eurelings, L. S., de Jonghe, J. F., Kalisvaart, K. J., Eikelenboom, P., & van

Gool, W. A. (2010). Delirium in elderly patients and the risk of postdischarge

mortality, institutionalization, and dementia: a meta-analysis. *JAMA*, 304(4), 443-451.

Yen, T. E., Allen, J. C., Rivelli, S. K., Patterson, S. C., Metcalf, M. R., Flink, B. J., . .

. Kwatra, M. M. (2016). Association between Serum IGF-I levels and Postoperative Delirium in Elderly Subjects Undergoing Elective Knee Arthroplasty. *Sci Rep*, 6, 20736.

Yudofsky, S. C., & Hales, R. E. (2008). *The American Psychiatric Publishing textbook of neuropsychiatry and behavioral neurosciences* (5th ed.). Washington, DC: American Psychiatric Pub.

Table 1

Summarization of recruited studies in current meta-analysis

Author (year)	Criteria	Population or setting	Delirium		Source	IGF (ng/mL)	Country
			Mean age	Female (%)			
			Non-delirium				
Yen, T. E. (2016)	CAM + DSM-IV	Elective knee arthroplasty	22			57.0±18.6	
			73.4±5.1	52.0	serum		USA
			76			68.4±31.6	
Shen, H. (2016)	CAM + DSM-IV	Open abdominal surgery	36			50.4±16.2	
			70.1±6.4	57.2	serum		China
			104			67.0±24.1	
Chu, C. S. (2016)	CAM + DSM-IV	Orthopedic surgery	23			97.1±47.4	
			81.7±4.0	26.2	serum		Taiwan
			80			96.1±47.6	
Egberts, A.	DOSS + DSM-IV	Internal and geriatric ward	23	82.6	53.5 serum	*48.1±32.1	Netherland

(2015)			63			71.0±43.5
Cinar, M. A.	DSM-IV	Cardiovascular surgery	15	64.2±10.3	25.7 serum	1.44±1.76
(2014)			20			1.03±1.18
Cerejeira, J.	CAM + DSM-	Elective hip arthroplasty	37	73.0±6.3	50.5 plasma	*138.3±57.8
(2013)	IV		64			128.2±59.6
Morandi, A.	CAM + CAM-	Critically ill mechanically ventilated patients	86	65	48.2 serum	#n/a USA
(2011)	ICU		24			
Lemstra, A.	CAM + DSM-	Acute or elective hip surgery	18	78.9	69.1 blood	*109.9
W.	IV		50			98.4
(2008)						
Adamis, D.	CAM	Elderly medical unit	22	84.6±6.6	67.1 serum	55.8±33.7
(2007)			35			75.2±33.7

Wilson, K. (2005)	CAM + DSM- III	Acute medical ward	12 88	84.5±4.2	69.0 plasma	[#] n/a UK
----------------------	-------------------	--------------------	----------	----------	-------------	---------------------

*: nmol/L convert to ng/ml x 7.63

[#]: derived effect sizes from other statistical parameters

Abbreviation: CAM: confusion assessment method; DOSS: Delirium observation screening scale; ELISA: enzyme-linked immunosorbent assay; IGF: insulin-like growth factor; n/a: not available; RASS: Richmond agitation-sedation scale; RIA: radioimmunoassay; UK: United Kingdom; USA: United States of America.

Figure Caption

Table 1

Summarization of recruited studies in current meta-analysis

Author (year)	Criteria	Population or setting	Delirium Non-delirium	Mean age	Female (%)	Source	IGF (ng/mL)	Country
Yen, T. E. (2016)	CAM + DSM-IV	Elective knee arthroplasty	22 76	73.4±5.1	52.0	serum	57.0±18.6 68.4±31.6	USA
Shen, H. (2016)	CAM + DSM-IV	Open abdominal surgery	36 104	70.1±6.4	57.2	serum	50.4±16.2 67.0±24.1	China
Chu, C. S. (2016)	CAM + DSM-IV	Orthopedic surgery	23 80	81.7±4.0	26.2	serum	97.1±47.4 96.1±47.6	Taiwan
Egberts, A. (2015)	DOSS + DSM-IV	Internal and geriatric ward	23 63	82.6	53.5	serum	*48.1±32.1 71.0±43.5	Netherland
Cinar, M. A. (2014)	DSM-IV	Cardiovascular surgery	15 20	64.2±10.3	25.7	serum	1.44±1.76 1.03±1.18	Turkey
Cerejeira, J. (2013)	CAM + DSM-IV	Elective hip arthroplasty	37 64	73.0±6.3	50.5	plasma	*138.3±57.8 128.2±59.6	Portugal
Morandi, A. (2011)	CAM + CAM-ICU	Critically ill mechanically ventilated patients	86 24	65	48.2	serum	#n/a	USA

Lemstra, A. W. (2008)	CAM + DSM- IV	Acute or elective hip surgery	18 50	78.9	69.1 blood	*109.9 98.4	Netherland
Adamis, D. (2007)	CAM	Elderly medical unit	22 35	84.6±6.6	67.1 serum	55.8±33.7 75.2±33.7	UK
Wilson, K. (2005)	CAM + DSM- III	Acute medical ward	12 88	84.5±4.2	69.0 plasma	#n/a	UK

*: nmol/L convert to ng/ml x 7.63

#: derived effect sizes from other statistical parameters

Abbreviation: CAM: confusion assessment method; DOSS: Delirium observation screening scale; ELISA: enzyme-linked immunosorbent assay; IGF: insulin-like growth factor; n/a: not available; RASS: Richmond agitation-sedation scale; RIA: radioimmunoassay; UK: United Kingdom; USA: United States of America.


Figure legends:**Figure 1.** Flow chart showing the selection strategy and inclusion/exclusion criteria employed in our current meta-analysis.

Figure 2. Forest plot of results of pooled effect sizes (Hedges' g) and 95% confidence intervals (CIs) from pooled results of peripheral levels of insulin growth factor-1 (IGF-1) between delirious and non-delirious patients within the entire population.




Figure 2 Forest plot of results of pooled effect sizes of recruited studies